

## DAFTAR PUSTAKA

- AOAC. (2005). Official Methods of Analysis. 18th ed. Association of Official Analytical Chemists; Arlington, VA, USA.
- AOAC. (1984). Official Methods of Analysis. 11<sup>th</sup> edition. Association of Official Analytical Chemists Inc., Washington, D.C
- Ardiansyah. (2016). "Mengenal bekatul lebih jauh". Program Studi Ilmu dan Teknologi Pangan, FTIK, Universitas Bakrie. [Online]. <http://www.bakrie.ac.id/id/berita-itp/artikel-pangan/929-mengenal-bekatul-lebih-jauh>. Diakses pada tanggal 13 Februari 2018.
- Ardiansyah., Gustriani, D., Handoko, D., Kusbiantoro, B., and Budijanto, S. "The antioxidant activity of indonesian fermented rice bran with Rhizopus oligosporus," in *Proceedings of the 15<sup>th</sup> ASEAN Conference on Food Science and Technology*. Vietnam. 463-468, Nov. 14-17, 2017.
- Ardiansyah., Shirakawa, H., Koseki, T., Ohinata, K., Hashizume, K., and Komai, M. (2006). Rice bran fractions improve blood pressure, lipid profile, and glucose metabolism in stroke-prone spontaneously hypertensive rats. *Journal of Agriculture and Food Chemistry*, 54: 1914-1920.
- Badan Litbang Pertanian. (2016). Pengaruh Lama Penyosohan terhadap Komposisi Kimia Beras. [Online] <http://www.litbang.pertanian.go.id/berita/one/2624/>. Diakses pada tanggal 11 Maret 2018
- Balai Pengelola Alih Teknologi Pertanian (BPATP). (2014). [Online] <http://bpatp.litbang.pertanian.go.id/ind/index.php/61-berita/864-beras-aromatik>. Diakses pada tanggal 11 Maret 2018
- Balai Pengkajian Teknologi Pertanian Bali. (2015). Pemanfaatan Bekatul, Limbah Penggilingan Padi sebagai Tepung Rendah Lemak. Bali. [Online] <http://bali.litbang.pertanian.go.id/ind/index.php/info-teknologi/567-pemanfaatan-bekatul-limbah-penggilingan-padi-sebagai-tepung-rendah-lemak>. Diakses pada tanggal 25 Januari 2018.
- Bobo-Garcia., G, Daidov-Pardo., G, Arroqui., C, Virseda P, Marin-Arroyo., MR, and Navarro M. (2015). Intra-laboratory validation of microplate methods for total phenolic content and antioxidant activity on polyphenolic extracts, and comparison with conventional spectrophotometric methods. J Sci Food Agric. 95(1): 204-209.
- Carlos, H.P., J.G. Chaves, R.R.S. Mundo, J. Namiesnik, S. Gorinstein, and G.A.G-Aguilar. (2012). Antioxidant interaction between major phenolics compounds

- found in 'Ataulfo' mango pulp: chlorogenic, gallic, orotocatechuic and vanillic acids. *Molecules*, 17: 12657-12664.
- Craft, BC., Kerrihard., AL, Amarowicz., R, and Pegg RB. (2012). Phenol-based antioxidants and the in vitro methods used for their assessment. *Compr Rev Food Sci Food Saf*. 11: 148-173.
- Dewi, N. M. A. P., Suter, I. K., & Widarta, I. W. R. (2012). Stabilisasi bekatul dalam upaya pemanfaatannya sebagai pangan fungsional. *Jurnal Ilmu dan Teknologi Pangan*.
- Endrawati, D., & Kusumaningtyas, E. (2018). Several Functions of Rhizopus sp on Increasing Nutritional Value of Feed Ingredient. *WARTAZOA. Indonesian Bulletin of Animal and Veterinary Sciences*, 27(2), 81-88.
- Forster GM, Raina K, Kumar A, Kumar S, Agarwal R, Chen MH, Bauer JE, McClung AM, Ryan EP. (2013). Rice varietal differences in bioactive bran components for inhibition of colorectal cancer cell growth. *Food Chem*. 141: 1545-1552
- Gustriani D. (2016). Perubahan total fenol dan aktivitas antioksidan bekatul (Sintanur dan Inpari 24) selama fermentasi dengan *Rhizopus oligosporus*. [Skripsi]. Jakarta (ID): Universitas Bakrie.
- Hartati S, Marsono Y, Suparmo, Santoso U. (2015). Komposisi kimia serta aktivitas antioksidan ekstrak hidrofilik bekatul beberapa varietas padi. *Agritech*. 35 (1): 35-42.
- Inagaki, S., Kato, T., Mori, S., & Fujita, T. (2013). Composition and Antioxidant Activity of Rice Fermented with Saccharifying Organisms from Asian Countries. *Food Science and Technology Research*, 19(5): 893-899.
- Islam, S., Nagasaka, R., Ohara, K., Hosoya, T., Ozaki, H., Ushio, H., & Hori, M. (2011). Biological abilities of rice bran-derived antioxidant phytochemicals for medical therapy. *Current Topics in Medicinal Chemistry*, 11(14): 1847-1853.
- Jung, T. D., Shin, G. H., Kim, J. M., Choi, S. I., Lee, J. H., Lee, S. J., Park, S. J., Woo, K.S., Oh, S.K, and Lee, O. H. (2017). Comparative analysis of  $\gamma$ -Oryzanol,  $\beta$ -Glucan, total phenolic content and antioxidant activity in fermented rice bran of different varieties. *Nutrients*, 9(6): 571.
- Kementrian Pertanian. (2017). Data Kementan Selaras Dengan Data BPS. Kementrian Pertanian. Jakarta. [Online] [http://www.pertanian.go.id/ap\\_posts/detil/1181/2017/09/28/09/30/05/Data%20Kementan%20Selaras%20Dengan%20Data%20BPS](http://www.pertanian.go.id/ap_posts/detil/1181/2017/09/28/09/30/05/Data%20Kementan%20Selaras%20Dengan%20Data%20BPS). Diakses pada tanggal 25 Januari 2018.

- Kupski, L. C.-S.-F.-s.-S. (2014). Solid-State Fermentation for the Enrichment and Extraction of Proteins and Antioxidant Compounds in Rice Bran by *Rhizopus oryzae*. *Braz. Arch. Biol. Technol.*, 55(6): 937.
- Kurniati, Y., Budijanto, S., Nuraida, L., Dewi, F. N. A. (2017). Peningkatan Senyawa Fenolik Bekatul dengan SSF (*Solid State Fermentation*) sebagai Pencegah Kanker. *Iptek Tanaman Pangan*, 12(2): 97-103.
- Kusumaningati, R.W. (2009). Analisis Kandungan Total Fenol Jahe (*Zingiber officinale Roscoe*) secara In Vitro. [Skripsi]. Jakarta: Fakultas Kedokteran, Universitas Indonesia.
- Lateef, A., Oloke, J. K., Kana, E. G., Oyeniyi, S. O., Onifade, O. R., Oyeleye, A. O., Oladosu, O.C and Oyelami, A. O. (2008). Improving the quality of agro-wastes by solid-state fermentation: enhanced antioxidant activities and nutritional qualities. *World Journal of Microbiology and Biotechnology*, 24(10): 2369-2374.
- Laokuldilok T, Surawang S, Klinhom J. 2013. Influence of miling time on the nutritional composition and antioxidant content of Thai rice bran. *Food and Applied Bioscience Journal*. 1 (3): 112-130.
- Magalhaes, L.M., F. Santos, M.A. Segundo, S. Reis, and J.L. Lima. (2010) Rapid microplate high-troughput methodology for assesment of *Folin-Ciocalteu* reducing capacity. *Talanta*. 83(2): 441-447.
- Min B, Gu L, McClung AM, Bergman CJ, Chen MH. (2012). Free and bound total phenolic concentrations, antioxidant capacities, and profiles of proanthocyanidins and anthocyanins in whole grain rice (*Oryza sativa* L.) of different bran colours. *Food Chem.* 133: 715-722.
- Moongngarm, A., Daomukda, N., and Khumpika, S. (2012). Chemical compositions, phytochemicals, and antioxidant capacity of rice bran, rice bran layer, and rice germ. *Journal of APCBEC Procedia*, 2: 73-79.
- Muntana, N., and Prasong, S. (2010). Study on total phenolic contents and their antioxidant activities of Thai white, red and black rice bran extracts. *Pakistan Journal of Biological Sciences*, 13(4): 170.
- Murtini, E. S.-A. (2016). Pengaruh Metode Fermentasi Substrat Padat dan Substrat Terendam pada Biji Sorghum terhadap Kualitas Tepung. *Jurnal Teknologi dan Industri Pangan*, 59-67.
- Nagarajan, S., Jagadish, S. V. K., Prasad, A. H., Thomar, A. K., Anand, A., Pal, M., and Agarwal, P. K. (2010). Local climate affects growth, yield and grain quality of aromatic and non-aromatic rice in northwestern India. *Agriculture, Ecosystems & Environment*, 138(3-4): 274-281.

- Nema R K, Sulekha M, Satish Y, Sunita Y. (2009). Antioxidant : a review. *J Chem Pharm Res.* 1 (1): 102-104
- Oliveira, M. S., Cipolatti, E. P., Furlong, E. B., dan Soares, L. S. (2012). Phenolic compounds and antioxidant activity in fermented rice (*Oryza sativa*) bran. *Ciênc. Tecnol. Aliment., Campinas*, 32(3): 531-537.
- Oliveira, M. S., Federn, V., Kupski, L., Cipolatti, E.P., Badiale-Furlong, E., and Souza-Soares, L.A. (2010). Physicochemical characterization of fermented rice bran biomass. *Journal of Food*, 8: 229–236.
- Razak, D. L. A., Rashid, N. Y. A., Jamaluddin, A., Sharifdin, S. A., and Long, K. (2014). Enhancement of phenolic acid content and antioxidant activity of rice bran fermented with *Rhizopus oligosporus* and *Monascus purpureus*. *Biocatalysis and Agricultural Biotechnology*, 4(1): 33-38.
- Razak, D. L. A., Rashid, N. Y. A., Jamaluddin, A., Sharifudin, S. A., Kahar, A. A., and Long, K. (2017). Cosmeceutical potentials and bioactive compounds of rice bran fermented with single and MIX culture of *Aspergillus oryzae* and *Rhizopus oryzae*. *Journal of the Saudi Society of Agricultural Sciences*, 16(2): 127-134.
- Ryan, E. P., Heuberger, A. L., Weir, T. L., Barnett, B., Broeckling, C. D., and Prenni, J. E. (2011). Rice bran fermented with *Saccharomyces boulardii* generates novel metabolite profiles with bioactivity. *Journal of Agriculture Food Chemistry*, 59(18): 1862-1870.
- Saidah., Syafruddin, dan Pangestuti R. (2015). Daya hasil padi sawah varietas Inpari 24 di beberapa lokasi SL-PTT di Sulawesi Tengah. *Pros Sem Nas Masy Biodiv Indon*, 1 (5): 1147-1150.
- SÁNCHEZ, C. (2009). Lignocellulosic residues: Biodegradation and bioconversion by fungi. *Biotechnology Advances*, 27: 185-194. PMid:19100826. <http://dx.doi.org/10.1016/j.biotechadv.2008.11.001>
- Schmidt, C. G., Gonçalvesa, L. M., Priettoa, L., Hackbart, H. S., and Furlong, E. B. (2014). Antioxidant activity and enzyme inhibition of phenolic acids from fermented rice bran with fungus *Rhizopus oryzae*. *Food Chemistry*, 146, 371-377.
- Sompong, R., Siebenhandl-Ehn, S., Linsberger-Martin, G., and Berghofer, E. (2011). Physicochemical and antioxidative properties of red and black rice varieties from Thailand, China and Sri Lanka. *Food Chemistry*, 124(1): 132-140.

- Sukma, L. N., Zackiyah., dan Gumilar. G. G. (2010). Pengkayaan asam lemak tak jenuh pada bekatul dengan cara fermentasi padat menggunakan *Aspergillus terreus*. *Jurnal Sains dan Teknologi Kimia*, 1(1), 66-72.
- Suprihatin. (2010). Teknologi Fermentasi. Surabaya: UNESA University Press.
- Susanti, Hari dan R. Alfian. (2012). Penetapan Kadar Fenolik Total Ekstrak Metanol Kelopak Bunga Rosella Merah (*Hibiscus sabdariffa* Linn) dengan Varietas Tempat Tumbuh Secara Spektrofotometri. *Jurnal Ilmiah Kefarmasian.*, 2(1): 73-80.
- Swastika, N.D. (2009). Stabilisasi Tepung Bekatul melalui Metode Pengukusan dan pengeringan Rak serta Pendugaan Umur Simpannya. [Skripsi]. Jurusan Teknologi Industri Pertanian, IPB.
- Tabaraki, R., & Nateghi, A. (2011). Optimization of ultrasonic-assisted extraction of natural antioxidants from rice bran using response surface methodology. *Ultrasonics sonochemistry*, 18(6): 1279-1286.
- Tarigan, E. B., dan Kusbiantoro, B. (2015). Pengaruh Derajat Sosoh dan Pengemas terhadap Mutu Beras Aromatik selama Penyimpanan. *Jurnal Penelitian Pertanian Tanaman Pangan*, 30(1).
- Wahab, M.I., Satoto., Rachmat, R., Guswara, A., dan Suwarna. (2017). Deskripsi varietas unggul baru padi. Badan Penelitian dan Pengembangan Pertanian. Subang, 27.
- Wijaya, H., B. Kusbiantoro, D.N. Faridah, D.D. Handoko, dan Taufik. (2008). Identifikasi komponen aroma aktif beberapa varietas beras (*Oryza miristica L*) aromatik asli Indonesia sebagai upaya pemanfaatan potensi beras Indonesia. Kerjasama Kemitraan Penelitian Pertanian dengan Perguruan Tinggi (KKP3T), Departemen Pertanian. Jakarta.
- Wulandari AP, Suter KI, Putra KN, Widarta RIW. (2012). Bekatul beras merah sebagai salah satu alternatif sumber antioksidan. *J Itepa*. 1 (1): 1-8.
- Yawadio, R., Tanimori, S., dan Morita, N. (2007). Identification of phenolic compounds isolated from pigmented rices and their aldose reductase inhibitory activities. *Food Chemistry*, 101(4): 1616-1625.
- Zubaidah, E., Nawa A., dan Fitri C.N. (2010). Studi aktivitas antioksidan bekatul dan susu bubuk skim terfermentasi bakteri asam laktat probiotik (*Lactobacillus plantarum* J2 dan *Lactobacillus casei*). *Jurnal Teknologi Pertanian*, 11(1): 11-17.

Zubaidah E dan Dinson DP. (2015). Pembuatan kulit pizza bekatul (kajian perlakuan stabilisasi dan proporsi tepung bekatul: tepung terigu). *JPA*, 3(1): 32-40.